



Science Bulletin

Fish Farm Link to Sea Lice Infections on B.C. Wild Salmon Confirmed

Transmission Dynamics of Parasitic Sea Lice from Farm to Wild Salmon


Martin Krkošek

Centre for Mathematical Biology, Depts. of Mathematical and Statistical Sciences and Biological Sciences
University of Alberta, Edmonton, AB

Salmon farming has been associated with sea lice infestations of wild juvenile salmon and wild salmon population declines in B.C.'s Broughton Archipelago. Despite a strong weight of evidence, the causal nature of these relationships has been highly contentious. A scientific consensus has been limited, in part, by an inability to adequately track the movement of these parasites between farmed and wild salmon.

Infection pressure near the farm was approximately 70 times greater than natural levels & exceeded ambient levels for 30 km.

Research Summary This peer-reviewed report looks at lice infestations on wild juvenile pink and chum salmon as they migrated past an isolated salmon farm down a long and narrow migration corridor in the Broughton Archipelago, BC (see Box 1). Mathematical models were used to analyze these data (see Box 2) and revealed juvenile salmon were initially infected with sea lice originating from two host populations: farm salmon and naturally occurring hosts. The calculations suggest the infection pressure near the farm was approximately 70 times greater than natural ambient levels and exceeded ambient levels for 30 km of migration route (see Box 3). This amounts to a total direct contribution of sea lice from the farm that was approximately 30,000 times greater than the natural production of sea lice in the length of habitat occupied by the salmon farm.



Box 1. Study Area

- ★ Sample sites,
- Active salmon farms

Salmon Migration Juvenile pink and chum salmon originate in freshwater streams in Knight Inlet. They must then travel south and west down the inlet toward Tribune channel and either continue down Knight Inlet or travel through Tribune Channel on their way to sea.

Sample Sites Farm A is the first salmon farm juvenile salmon would encounter along their migration and is the focus of this analysis presented here.


Data All sites were sampled within a 14-day period in May 2003 resulting in a high-resolution dataset of sea lice infection levels on juvenile pink and chum salmon as they passed an isolated salmon farm (Farm A). The non-lethal sampling of approximately 5500 fish far surpasses the sampling intensity other B.C. studies. Please refer to the source paper^{iv} and Krkošek et al 2005b^v for the full methodology and analysis.

Second Generation of Farm-Origin Lice Causes

Additional Infection The results also show that lice, once transmitted to wild juvenile salmon, were transported down the migration route where they reproduced and re-infected the wild juvenile salmon. Inclusion of this second generation of lice raises the farm-origin infection pressure above natural levels for approximately 75 km of migration route resulting in a total direct and indirect contribution of sea lice that was 200,000 times greater than the natural production of sea lice in the length of habitat occupied by the salmon farm. This transport and reproduction of farm-origin lice raises the possibility for disease growth and spread to other wild salmon populations up and down the coast.

These data satisfy even the most conservative benchmark for proof - this is the definitive work on the issue.

Box 2. Louse Lifecycle and Transmission Dynamics





Sea Lice (*Lepeophtheirus salmonis* and *Caligus clemensi*)

Sea lice have two distinct phases in their lifecycles – planktonic and parasitic. Planktonic larvae float freely in the water and allow lice to be transmitted between wild and farmed salmon. During the second, parasitic phase, lice attach themselves to a host salmon and feed on the surface of the fish – leading to increased disease and sometimes death in their infected hosts (See Box 2).

Links to Salmon Farms

Several European studies have correlated salmon farming with sea lice infestations of wild fishⁱ. Several studies in B.C have examined this correlation and one study found that wild pink juvenile salmon were more heavily infected in salmon farming areas than in areas with no salmon farmsⁱⁱ. An additional study analyzed these relationships and discounted other non-salmon farm variablesⁱⁱⁱ. The study described here is the first to quantify the spatial footprint of sea lice transmission from farm to wild salmon and track the subsequent farm-origin lineage of lice that infected wild salmon^{iv}.

These data, due to the massive sampling effort and the unequivocal nature of the conclusions, satisfy even the most conservative benchmark for proof - this is the definitive work on the issue. Sea lice counts released by Stolt Sea Farms confirm that study captures an average farm with sea lice numbers and stocking densities well within industry norms. Therefore the resultant transmission data are to be viewed as a conservative estimate of the impact of an “average” farm in the Broughton Archipelago.



Box 3. Model Agrees With Field Data

Graph (b) shows the model’s inferred origins and spatial profiles of sea lice larvae that would be required to produce the patterns seen in the data (a). The red lines depict the location and relative abundances of planktonic larvae whose origin can be traced back to farmed salmon either directly or through one generation of lice.

Panels (a), above on the left, show actual louse abundances on juvenile pink and chum salmon as they passed Farm A, located at $x=0$. Points with 95% confidence bounds correspond to field data and the solid lines are the best fit of the model to the data.

Science Bulletin

Fish Farm Link to Sea Lice

Infections on B.C. Wild Salmon Confirmed

Implications

Given the increasing weight of evidence, including this research^{iv}, that salmon farms are a potentially lethal source of sea lice on wild salmon, there is ample evidence to compel precautionary action by regulators. The premise of industrial-scale open net aquaculture in wild salmon habitats needs to be reconsidered. The proposed expansion of the industry, including the introduction of new farmed species, throughout BC must be halted until the full ecological costs of this industry are understood. There is a clear potential for severe and irreversible damages to be inflicted upon wild salmon populations and their dependent cultures, ecosystems, and economies.

Research Published in Peer Reviewed Journal

The paper described here is published in the Proceedings of the Royal Society of London, Series B, March 2005 as, Krkošek, M., Lewis, M. A., & Volpe, J. P. *Transmission dynamics of parasitic sea lice from farm to wild salmon.*

Acknowledgements

The David Suzuki Foundation, Raincoast Conservation Society, Raincoast Research Society, several National Science and Engineering Research Council (NSERC) grants, a Canada Research Chair and a Walter H. Johns Graduate Fellowship, supported this work. David Suzuki Foundation was Industrial Partner for M. Krkošek's NSERC Scholarship.



Salmon fry with sea lice
A. Morton photo

ⁱMcVicar, A. H. 2004 Management actions in relation to the controversy about salmon lice infections in fish farms as a hazard to wild salmonid populations. *Aquaculture Research*. 35, 751–758.

ⁱⁱ Morton, A., Routledge, R., Peet, C., & Ladwig, A. 2004 Sea lice (*Lepeophtheirus salmonis*) infection rates on juvenile pink (*Oncorhynchus gorbuscha*) and chum (*O. Keta*) salmon in the near shore marine environment of British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences*. 61,147-157.

ⁱⁱⁱ Williams, I., Groot, C., & Walthers, L. 2004. Possible Factors Contributing to the Low Productivity of the 2000 brood Year Pink Salmon (*Oncorhynchus gorbuscha*). *David Suzuki Foundation*, 2004.

^{iv} Krkošek, M., Lewis, M. A., & Volpe, J. P. 2005a Transmission dynamics of parasitic sea lice from farm to wild salmon. *Proceedings of the Royal Society of London Series B*, in press; on-line 30 March 2005.

^v Krkošek, M., Morton, A., & Volpe, J. P. 2005b Non-lethal assessment of juvenile pink and chum salmon for parasitic sea lice infections and fish health. *Transactions of the American Fisheries Society*, in press.



David
Suzuki
Foundation

